



## The role of volatile organic compounds in the polluted urban atmosphere of Bristol, UK

A. C. Rivett, D. Martin, D. J. Gray, C. S. Price, G. Nickless, P. G. Simmonds, S. J. O'Doherty, B. R. Grealley, A. Knights, D. E. Shallcross

### ► To cite this version:

A. C. Rivett, D. Martin, D. J. Gray, C. S. Price, G. Nickless, et al.. The role of volatile organic compounds in the polluted urban atmosphere of Bristol, UK. *Atmospheric Chemistry and Physics Discussions*, 2003, 3 (1), pp.769-796. hal-00300926

**HAL Id: hal-00300926**

**<https://hal.science/hal-00300926>**

Submitted on 13 Feb 2003

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**VOCs in the polluted  
urban environment of  
Bristol**

A. C. Rivett et al.

# The role of volatile organic compounds in the polluted urban atmosphere of Bristol, UK

**A. C. Rivett, D. Martin, D. J. Gray, C. S. Price, G. Nickless, P. G. Simmonds, S. J. O'Doherty, B. R. Grealley, A. Knights, and D. E. Shallcross**

Biogeochemistry Research Centre, School of Chemistry, University of Bristol, Cantock's Close, Bristol, BS8 1TS, UK

Received: 15 August 2002 – Accepted: 29 January 2003 – Published: 13 February 2003

Correspondence to: A. C. Rivett (a.c.rivett@bristol.ac.uk)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

## Abstract

The results of a field campaign carried out from early spring through to the late summer of 2000, in Bristol, England, are presented. Continuous measurements of over 40 hydrocarbons have been made at an urban background site, located at Bristol University, for approximately nine months using a GC-FID system and for a selection of halocarbons for approximately one month using a GC-ECD system. In this paper we present the time-series of the nine halocarbons and selected hydrocarbons. Daytime and night-time hydroxyl radical concentrations have been estimated based on the diurnal variations of a selection of the measured hydrocarbons. The average summer daytime concentration of OH was found to be  $2.5 \times 10^6$  molecules  $\text{cm}^{-3}$  and the night-time concentration to be in the range  $10^4$  to  $10^5$  molecules  $\text{cm}^{-3}$ . In addition, the role played by certain VOCs in the formation of ozone is assessed using the POCP (Photochemical Ozone Creation Potential) concept.

## 1. Introduction

The urban environment contains a complex mixture of chemicals, however, due to the high levels of  $\text{NO}_x$  that generally persist, ozone formation is VOC (volatile organic compound) limited (e.g. Sillman, 1999). Therefore, it is of great importance to determine the type of VOC that is present in the urban environment, its concentration and how this varies both spatially and temporally. This study has investigated both local and remote effects on levels of pollutants in the Bristol conurbation. A detailed understanding of the different influences on local air quality is important to inform relevant policy decisions such as transport planning and healthcare provision and this type of research provides such information.

The behaviour of halocarbons in the stratosphere and the remote troposphere is well understood due to their major role in ozone depletion, yet there have been few studies in urban areas. A handful of investigations have been carried out in Taiwan, Brazil

ACPD

3, 769–796, 2003

## VOCs in the polluted urban environment of Bristol

A. C. Rivett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

and Japan (Wang et al., 1998; Grosjean et al., 1999; Yamamoto et al., 2000). There are several reasons for the study halocarbons in urban atmospheres. Their degradation products are sources of the Cl radical which acts like the OH radical and so may change the oxidising capacity of the troposphere. In addition, the health effects are significant, these compounds are potentially toxic, irritant and carcinogenic. Emissions in Britain are regulated by the UK Environment Agency, but are not systematically monitored.

In urban conurbations, the sources and behaviour of hydrocarbons are generally well identified and understood (Wayne, 2000; UK PORC, 1997; Derwent et al., 2000). Their overwhelming source is rush-hour traffic, inducing a diurnal cycle upon which is added transported air masses bringing polluted or clean air from other regions. In addition to being a measure of air quality, the hydrocarbon measurements can also provide useful information about the chemistry of the polluted atmosphere. For example, the formation of ozone and other photochemical pollutants in the NO<sub>x</sub>-rich urban atmosphere is VOC-limited, so it is useful to determine which species are making the greatest contribution to ozone formation. Concentrations of OH and other radicals can be calculated during the day and night, as can the magnitude of dispersion processes taking place. Simple source identification is also possible using these types of data but this analysis is not discussed here.

## 2. Experimental

Bristol (51.5° N, 2.75° W), with a population of almost half a million people, is situated in the south-west of Britain, 120 miles west of London. Like any other town or city, Bristol has its own distinctive atmospheric characteristics. It is located a short way from the coast, and is surrounded by hills. The city centre basin covers an area of approximately 25 km<sup>2</sup>, although the suburbs extend much further. To the west of the city is a large industrial area (Avonmouth) and other key industries include manufacturing and business. Two major motorways run nearby and the volume of traffic around the area is generally high. The basin geography combined with heavy traffic flow can lead to

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Print Version	
Interactive Discussion	

the build-up of pollutants during anti-cyclonic weather conditions, although prevailing winds are often from the clean, western Atlantic sector.

From early spring to late summer 2000, continuous measurements of VOCs were made at an urban background site, located at the University of Bristol. The Automatic Adsorption-Desorption System (ADS) and GC technology developed at Bristol (Simmonds et al., 1995; Sheppard, 1999; Bassford et al., 1998) was used to perform routine analysis of air and standard samples in a continuous cycle. Over 40 hydrocarbon species were measured using a GC-FID system with a Peltier-cooled ADS to achieve low temperature trapping at minus 40° C over a period of approximately 9 months. A GC-ECD was deployed concurrently at the same sampling point to monitor a selection of 9 halocarbons for one month using an ADS which traps at ambient temperature (Bassford et al., 1998). Measurements of a smaller set of hydrocarbons were taken simultaneously at a roadside site in the centre of Bristol, as part of the UK national monitoring network, NETCEN (Derwent et al., 2000). The species monitored by the three systems were; C<sub>2</sub>–C<sub>8</sub> Hydrocarbons, CFCs 11 and 12, Halons 1211 and 1301, CHCl<sub>3</sub> (Chloroform), CH<sub>3</sub>CCl<sub>3</sub> (Methyl Chloroform), CCl<sub>4</sub> (Carbon Tetrachloride), TCE (Trichloroethene) and PCE (Perchloroethene).

### 3. Halocarbons in the city

Figures 1 to 3 show the concentrations of the nine halocarbons measured in Bristol between August and September 2000. A complex time-series is observed for the halocarbons, with large variations in the concentrations measured. There are some episodes where several compounds are simultaneously elevated, however there are also periods where just one compound is elevated, in complete contrast to the coherent variations observed for most hydrocarbons, which are driven primarily by vehicular sources. Figure 4 shows the halocarbon time-series overlaid with toluene as a representative hydrocarbon.

No weekly or diurnal trends were found for any of the halocarbons, as might be ex-

## VOCs in the polluted urban environment of Bristol

A. C. Rivett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

pected for compounds with no significant biogenic or automobile sources. In general, halocarbon levels measured at the urban site are similar to those measured at Mace Head, Ireland, a rural background site which is part of the AGAGE monitoring network. The only exceptions are chloroform and methyl chloroform which are significantly elevated above background levels. Results of the halocarbon monitoring campaign and correlations between the halocarbons and hydrocarbons are discussed further in Rivett et al. (2003).

#### 4. Hydrocarbons in the city

The summer time-series of a representative selection of the hydrocarbons measured are shown in Fig. 5. The concentration time-series' are very similar for each of the compounds measured. This is consistent with what would be expected as these compounds all have a primarily vehicular source. There are periods of high pollution when many compounds are elevated and "clean" periods where most species are at relatively low levels. In the sections which follow, the hydrocarbon dataset is interrogated to estimate both daytime and night-time OH levels.

In the analysis of this type of dataset, investigation of the variation in hydrocarbon ratios can be a powerful tool (Calvert, 1976). Calculating the ratios of each of the hydrocarbons to one reference compound enables differences in their sources and reactivity to be elucidated. Using ratios removes day-to-day variations, meteorological factors and differences in emission strengths between the compounds (Roberts et al., 1984; Parrish et al., 1998). In this study, benzene has been used as the reference compound because its lifetime is longest of all the compounds measured (it has the slowest reaction rate with OH) and no significant non-traffic sources have been characterised in the UK VOC emission inventory (Derwent et al., 2000). Therefore, as an air mass is chemically processed, the ratios of all the other species to benzene will decrease.

### VOCs in the polluted urban environment of Bristol

A. C. Rivett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

#### 4.1. Daytime OH

It is well established that the OH radical is the main initiator of atmospheric chemistry during the day (Wayne, 2000). The concentration of the hydroxyl radical during the day was estimated from the variation in concentration ratios of four aromatic species (ethyl benzene, toluene, *o*-xylene and *m* + *p*-xylene) to benzene (Blake et al., 1993; Moschonas et al., 2001). Aromatics are most suitable for this type of calculation because they have similar lifetimes and sources to benzene and decay almost exclusively by reaction with the hydroxyl radical. From the change in a compounds' ratio to benzene between the morning rush hour period (07:00–10:00) and the midday period (10:00–16:00) the concentration of daytime OH can be calculated using the following equation:

$$[OH] = \frac{\ln \left( \frac{X_t/B_t}{X_i/B_i} \right)}{t(K_B - K_X)} \quad (1)$$

$X_i/B_i$  is the ratio of hydrocarbon  $X$  to benzene ( $B$ ) during the morning rush hour,  $X_t/B_t$  is the ratio of hydrocarbon  $X$  to benzene ( $B$ ) over midday,  $t$  is the time elapsed between morning rush hour and midday,  $K_B$  is the rate constant for the reaction of benzene and OH and  $K_X$  is the rate constant for the reaction of hydrocarbon  $X$  with OH. This gives an average OH concentration over the time elapsed.

Ratios to benzene were calculated over the entire campaign to give a typical ratio for each of the four aromatic species during the morning rush hour and midday time-periods. These ratios were then used to determine an average daytime OH concentration during the summer. The mean of the four different OH concentrations calculated from the different aromatic species is given in Table 1 along with the maximum and minimum concentrations calculated.

The lifetime of OH is less than 1 second and consequently noontime concentrations generally only reach 0.04–0.4 ppt ( $10^6$ – $10^7$  molecules  $\text{cm}^{-3}$ ) (Wayne, 2000). The average OH concentration of  $2 \times 10^6$  molecules  $\text{cm}^{-3}$  calculated for the summer period

### VOCs in the polluted urban environment of Bristol

A. C. Rivett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

compares well with what would be expected. Field campaigns such as the EASE 1996 and 1997 European background air experiments (Creasey et al., 1997; Creasey et al., 2002) and BERLIOZ (Platt et al., 2002), sampling in the Berlin area, have measured daytime [OH] of between  $1\text{--}10 \times 10^6$  molecules  $\text{cm}^{-3}$  during summer-time.


#### 5 4.2. Night-time OH

Although  $\text{NO}_3$  is generally thought to be the main oxidising species at night, hydroxyl radicals can be generated even after dark from several reactions (Platt et al., 1990; Mihelcic et al., 1993; Canosa-Mas et al., 1996; Carslaw et al., 1997; Paulson et al., 1997).

10 OH can be formed from the reaction of  $\text{HO}_2$  with the nitrate radical (Eq. 2a) or with ozone (Eq. 2b):



15 The reaction of alkenes with the nitrate radical can lead to the formation of  $\text{HO}_2$ , which can be converted to OH via reactions (2a) and (2b):



The ozonolysis of alkenes is a complex reaction, believed to involve the formation of a Criegee intermediate (Eqs. 4 and 5). However, strong evidence exists to suggest that

20 OH can be formed from these reactions which can proceed all day:



25 [OH] can be calculated from an analysis of the decay of VOCs during the night in 2 ways:

## VOCs in the polluted urban environment of Bristol

A. C. Rivett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion



#### 4.2.1. Alkene decays

Data from nights (such as that shown in Fig. 6) where several alkene species undergo a steady decay due to dispersion and reaction with OH, NO<sub>3</sub> and O<sub>3</sub> has been compiled. The rates of reaction of the VOCs and the VOC concentration over the night are known, therefore re-arranging the equation below, radical concentrations can be calculated.

$$d[\text{VOC}]/dt = (-k[\text{OH}] - k[\text{NO}_3] - k[\text{O}_3] - k_{\text{dispersion}}) \cdot [\text{VOC}] \quad (6)$$

A matrix method was used to derive the equations given in the appendix. The degradation of alkenes by ozone is ignored, as it is very slow in comparison with the reaction of alkenes with OH and NO<sub>3</sub>. Thus, decay data for 3 different alkenes is required on each night to perform these calculations. During the summer period of the campaign there were 7 such nights where sufficient data was collected.

The average night-time OH concentration was calculated to be  $3.8 \times 10^5$  molecules cm<sup>-3</sup>. The estimated OH concentrations varied over a range of  $6.3 \times 10^4$  molecules cm<sup>-3</sup> to  $1 \times 10^6$  molecules cm<sup>-3</sup>. Levels of NO<sub>3</sub> calculated by this method range from 2 to 7 ppt, with an average from the 7 nights being 4.2 ppt. This is consistent with (although somewhat lower than) NO<sub>3</sub> concentrations previously measured in urban areas (Platt et al., 2002; Smith et al., 1995). The concentration of NO<sub>3</sub> at night is strongly dependent on the levels of alkenes, with which it readily reacts. So in the polluted urban environment, NO<sub>3</sub> production could be substantially suppressed.

#### 4.2.2. Steady-state concentration

The steady-state concentration over one whole night can also be estimated using the following equation:

$$d[\text{OH}]/dt = k[\text{O}_3][\text{alkenes}](\text{all summed}) + k[\text{HO}_2][\text{NO}_3] - k[\text{OH}][\text{NO}_2][\text{M}] - k[\text{OH}][\text{CO}] - k[\text{OH}][\text{VOC}] \quad (7)$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

The change in OH is assumed to be equal to the production from the reaction of alkenes with ozone (production from  $\text{HO}_2 + \text{NO}_3$  is ignored, because the concentrations of these radicals are not known), minus the loss from reactions with  $\text{NO}_2$ , CO and other VOCs present.

5 Figure 7 shows OH levels calculated for each night of the campaign for the University site and for the city centre site. It can be seen that agreement is generally very good between the two sites. Average [OH] over the summer campaign period at the University was calculated to be  $3.6 \times 10^4$  molecules  $\text{cm}^{-3}$ , ranging from  $4.9 \times 10^3$  to  $1.4 \times 10^5$  molecules  $\text{cm}^{-3}$ . This is a much lower level than that calculated by the decay  
10 rate method because only 5 simple alkene species are used in the calculation and production from  $\text{HO}_2$  is ignored. If more alkenes were used in the calculation, particularly if heavier and more reactive alkene species were considered, such as isoprene and other monoterpenes, the estimation would be more accurate.

Few measurements have been made of OH at night in similar environments but some  
15 comparisons can be made. During the MLOPEX campaign at a remote background site in Hawaii, night-time [OH] was found to be  $1 \times 10^5$  molecules  $\text{cm}^{-3}$  (Eisele et al., 1996; Cantrell et al., 1997). The PROHET campaign, which took place in a heavily forested area of Michigan measured [OH] as high as  $1 \times 10^6$  molecules  $\text{cm}^{-3}$  after dark (Faloona et al., 2001). A similar campaign in a forest in northern Greece (AEROBIC)  
20 also found that OH in the order of  $10^5$  molecules  $\text{cm}^{-3}$  persisted into the night (Creasey et al., 2001). The BERLIOZ experiment (Platt et al., 2002), measuring in and around the Berlin area, (the area which is probably most similar to the environment studied here) found levels of  $1.4 \times 10^5$  molecules  $\text{cm}^{-3}$ . The hydroxyl radical concentrations of  $10^4$  to  $10^5$  molecules  $\text{cm}^{-3}$  calculated here, even by a relatively simple method, are  
25 nevertheless in good agreement with those found in other environments. It is reassuring to note that by using the data derived for [OH],  $[\text{NO}_3]$  and dispersion from the decay method, the calculated decay of measured alkanes is totally consistent.

## VOCs in the polluted urban environment of Bristol

A. C. Rivett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

### 4.3. Ozone creation potential

Classifying a wide variety of VOCs in terms of their ability to form ozone is a complex problem, since those compounds which are oxidised quickly lead to significant ozone formation close to the point of emission, whereas those which are oxidised more slowly will lead to ozone being formed further away from the source of the compound, for example in rural areas. Using the concept of Photochemical Ozone Creation Potential (POCP), the role played by each of the hydrocarbons in the creation of ozone has been revealed. In this work  $\varepsilon^{\text{POCP}}$  (estimated POCP) values have been used (Jenkin, 1998), which use a mathematical formula (as shown in Eq. 8) to calculate POCP, rather than using a modelling approach:

$$\varepsilon^{\text{POCP}} = \alpha_1 \cdot \gamma_S \cdot \gamma_R^\beta (1 - \alpha_2 \cdot \eta_C) \quad (8)$$

$\alpha_1$ ,  $\alpha_2$  and  $\beta$  are constants,  $\eta_C$  is the carbon number of the VOC and  $\gamma_S$  and  $\gamma_R$  are the structure-based and reactivity-based ozone formation indices respectively. This method replicates well POCPs calculated from the modelling work and also rationalises the POCP values in terms of the intrinsic structural and reactive properties of the VOCs themselves.

$\gamma_R$  is a simple approach which defines the relative ozone formation potential of a series of VOCs based on how quickly they form ozone in the boundary layer.  $\gamma_R$  is based on the rate coefficient ( $k_{\text{OH}}$ ) per reactive bond ( $\eta_B$ ) (Jenkin, 1998). This is normalised to a value of 1.0 for ethene as shown in Eq. (9):

$$\gamma_R = (k_{\text{OH}}/\eta_B) \cdot (6/k_{\text{OH}}^{\text{ethene}}) \quad (9)$$

$\gamma_R$  gives a better indication of the impact of VOCs on ozone formation on a local scale, compared to POCP, which gives an indication of the overall regional scale formation. In calculating  $\varepsilon^{\text{POCP}}$  and  $\gamma_R$  all compounds were ratioed to propane.

Plots of the proportion of ozone formed by each class of hydrocarbon as a percentage of the total are shown in Fig. 8. Alkenes produce the greatest amount of ozone

## VOCs in the polluted urban environment of Bristol

A. C. Rivett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

(45% of total) closest to the source of emissions (as calculated using  $\gamma_R$ ) with aromatic species generating a further 30%. The  $\varepsilon^{\text{POCP}}$  values calculated for the campaign show that alkane and aromatic hydrocarbon species have the greatest effect on regional scale ozone production, accounting for approximately 50% and 30%, respectively. This ties in very well with what would be expected. Close to the point of emission, more reactive compounds such as alkenes will contribute most to ozone formation. As time goes on, slower reacting compounds such as the alkanes will become more important in producing ozone.

## 5. Conclusions

Simultaneous measurements of nine different halocarbon species and over forty hydrocarbon species have been made over one summer at an urban background site in a large British city. This unusual dataset has provided a unique opportunity to compare two classes of compounds which display very different behaviour in the urban atmosphere.

The lifetimes of the hydrocarbons measured are all in the order of days, whereas those of the halocarbons measured vary much more considerably from a few days to over a hundred years. Similarly the halocarbons originate from a much more diverse range of sources than do the hydrocarbons studied. Whereas the  $\text{C}_2\text{--C}_8$  hydrocarbons measured at this typical urban site are overwhelmingly emitted by traffic, the halocarbons are emitted from a wide variety of industrial activity, both big factories and smaller businesses, from landfill and other waste processing activity and from the natural environment. Such diversity in lifetime and sources will naturally mean that the behaviour of hydrocarbons compared to halocarbons is very different but it also means that the various halocarbon compounds will not behave in a similar way to each other either. Consequently the time-series of the nine halocarbons are not simple to interpret, they do not display any significant diurnal cycle and each compound has shown large excursions from baseline levels at different points during the campaign.

## VOCs in the polluted urban environment of Bristol

A. C. Rivett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

A detailed analysis of the hydrocarbon dataset has been made possible by the large number of species measured during this campaign. Radical concentrations have been calculated for both day and night and the estimated OH concentrations compare well with field measurements made in similar environments. The contribution of different types of hydrocarbons to the formation of ozone in the city centre and further afield has also been quantified.

This type of investigation is an example of how such datasets can be a powerful tool to explore the chemistry and behaviour of urban environments. At this time, when there are numerous schemes to reduce different types of emissions in cities and towns across the world, analyses such as this can have a valuable part to play in informing the decision-making process.

## Appendix

These equations were derived using a matrix method and used to calculate OH, NO<sub>3</sub> and dispersion at night from the decay of alkenes.

$$[\text{OH}] = 1/\det * \left( \text{Slope}^{\text{VOC}_1} \left( k_{\text{NO}_3}^{\text{VOC}_2} - k_{\text{NO}_3}^{\text{VOC}_3} \right) + \text{Slope}^{\text{VOC}_2} \left( k_{\text{NO}_3}^{\text{VOC}_3} - k_{\text{NO}_3}^{\text{VOC}_1} \right) + \text{Slope}^{\text{VOC}_3} \left( k_{\text{NO}_3}^{\text{VOC}_1} - k_{\text{NO}_3}^{\text{VOC}_2} \right) \right) \quad (\text{A1})$$

$$[\text{NO}_3] = 1/\det * \left( \text{Slope}^{\text{VOC}_1} \left( k_{\text{OH}}^{\text{VOC}_3} - k_{\text{OH}}^{\text{VOC}_2} \right) + \text{Slope}^{\text{VOC}_2} \left( k_{\text{OH}}^{\text{VOC}_1} - k_{\text{OH}}^{\text{VOC}_3} \right) + \text{Slope}^{\text{VOC}_3} \left( k_{\text{OH}}^{\text{VOC}_2} - k_{\text{OH}}^{\text{VOC}_1} \right) \right) \quad (\text{A2})$$

## VOCs in the polluted urban environment of Bristol

A. C. Rivett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

## VOCs in the polluted urban environment of Bristol

A. C. Rivett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

$$k_{\text{dispersion}} = 1/\text{det} * \left( \text{Slope}^{\text{VOC}_1} \left( k_{\text{OH}}^{\text{VOC}_2} \cdot k_{\text{NO}_3}^{\text{VOC}_3} - k_{\text{NO}_3}^{\text{VOC}_2} \cdot k_{\text{OH}}^{\text{VOC}_3} \right) \right. \\ \left. + \text{Slope}^{\text{VOC}_2} \left( k_{\text{NO}_3}^{\text{VOC}_1} \cdot k_{\text{OH}}^{\text{VOC}_3} - k_{\text{OH}}^{\text{VOC}_1} \cdot k_{\text{NO}_3}^{\text{VOC}_3} \right) \right. \\ \left. + \text{Slope}^{\text{VOC}_3} \left( k_{\text{OH}}^{\text{VOC}_1} \cdot k_{\text{NO}_3}^{\text{VOC}_2} - k_{\text{NO}_3}^{\text{VOC}_1} \cdot k_{\text{OH}}^{\text{VOC}_2} \right) \right) \quad (\text{A3})$$

$$\text{Det} = k_{\text{OH}}^{\text{VOC}_1} \left( k_{\text{NO}_3}^{\text{VOC}_2} - k_{\text{NO}_3}^{\text{VOC}_3} \right) - k_{\text{NO}_3}^{\text{VOC}_1} \left( k_{\text{OH}}^{\text{VOC}_2} - k_{\text{OH}}^{\text{VOC}_3} \right) \\ + \left( k_{\text{OH}}^{\text{VOC}_2} \cdot k_{\text{NO}_3}^{\text{VOC}_3} - k_{\text{NO}_3}^{\text{VOC}_2} \cdot k_{\text{OH}}^{\text{VOC}_3} \right) \quad (\text{A4})$$

*Acknowledgement.* A. C. Rivett, D. J. Gray and C. S. Price would like to thank NERC for studentships and D. Martin thanks Bristol University for a scholarship. We also thank A. Knights for technical assistance during the campaign.

## References

- Bassford, M., Simmonds, P. G., Nickless, G.: An automated system for near-real-time monitoring of trace atmospheric halocarbons, *Anal. Chem.*, 70, 958–965, 1998.
- Blake, N. J., Penkett, S. A., Clemitshaw, K. C., Anwyl, P., Lightman, P., Marsh, A. R. W., and Butcher, G.: Estimates of atmospheric hydroxyl radical concentrations from the observed decay of many reactive hydrocarbons in well-defined urban plumes, *J. Geophys. Res.*, 98, 2851–2864, 1993.
- Calvert, J. G.: Hydrocarbon involvement in photochemical smog formation in Los Angeles atmosphere, *Environ. Sci. Technol.*, 10, 256–262, 1976.
- Canosa-Mas, C. E., King, M. D., Lopez, R., Percival, C. J., Wayne, R. P., Shallcross, D. E., Pyle, J. A., and Daele, V. J.: Is the reaction between  $\text{CH}_3\text{C}(\text{O})\text{O}_2$  and  $\text{NO}_3$  important in the night-time troposphere?, *J. Chem. Soc. Faraday Trans.*, 92, 2211–2222, 1996.

- Cantrell, C. A., Shetter, R. E., Calvert, J. G., Eisele, F. L., and Tanner, D. J.: Some considerations of the origin of nighttime peroxy radicals observed in MLOPEX 2, *J. Geophys. Res.*, 102, 15899–15913, 1997.
- 5 Carlsaw, N., Carpenter, L. J., Plane, J. M. C., Allan, B. J., Burgess, R. A., Clemitshaw, K. C., Coe, H., and Penkett, S. A.: Simultaneous observations of nitrate and peroxy radicals in the marine boundary layer, *J. Geophys. Res.*, 102, 18917–18933, 1997.
- Creasey, D. J., Halford-Maw, P. A., Heard, D. E., Pilling, M. J., and Whitaker, B. J.: Implementation and initial deployment of a field instrument for measurement of OH and in the troposphere by laser-induced HO<sub>2</sub> fluorescence, *J. Chem. Soc., Faraday Trans.*, 93, 2907–2913, 10  
1997.
- Creasey, D. J., Heard, D. E., and Lee, J. D.: OH and HO<sub>2</sub> measurements in a forested region of north-western Greece, *Atmos. Environ.*, 35, 4713–4724, 2001.
- Creasey, D. J., Heard, D. E., and Lee, J. D.: Eastern Atlantic Spring Experiment 1997 (EASE97) 1. Measurements of OH and HO<sub>2</sub> concentrations at Mace Head, Ireland, *J. Geophys. Res.*, 15  
107, 10.1029/2001JD001568, 2002.
- Derwent, R. G., Davies, T. J., Delaney, M., Dollard, G. K., and Field, R. A.: Analysis and interpretation of the continuous hourly monitoring data for 26 C<sub>2</sub>–C<sub>8</sub> hydrocarbons at 12 UK sites during 1996, *Atmos. Environ.*, 34, 297–312, 2000.
- Eisele, F. L., Tanner, D. J., Cantrell, C. A., and Calvert, J. G.: Measurements and steady-state calculations of OH concentrations at Mauna Loa Observatory, *J. Geophys. Res.*, 101,  
20 14665–14679, 1996.
- Faloona, I., Tan, D., Brune, W., Hurst, J., and Barket Jr, D.: Nighttime observations of anomalously high levels of hydroxyl radicals above a deciduous forest canopy, *J. Geophys. Res.*, 106, 24315–24333, 2001.
- 25 Grosjean, E., Rasmussen, R. A., and Grosjean, D.: Toxic air contaminants in Porto Alegre, Brazil. *Environ. Sci. Technol.*, 33, 1970–1978, 1999.
- Jenkin, M. E.: Photochemical Ozone and PAN Creation Potentials: Rationalisation and Methods of Estimation. AEAT Report 4182/20150/003, Issue 1, National Environmental Technology Centre, UK, 1998.
- 30 Mihelcic, D., Klemp, D., Musgen, P., Patz, H. W., and Volz-Thomas, A.: Simultaneous measurements of peroxy and nitrate radicals at Schauinsland, *J. Atmos. Chem.*, 16, 4, 313–335, 1993.
- Moschonas, N., Glavas, S., and Kouimtzis, T.: C<sub>3</sub> to C<sub>9</sub> hydrocarbon measurements in the

## VOCs in the polluted urban environment of Bristol

A. C. Rivett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

two largest cities of Greece, Athens and Thessaloniki, Calculation of hydrocarbon emissions by species, Derivation of hydroxyl radical concentrations, Sci. Total Environ., 271, 117–133, 2001.

Parrish, D. D., Trainer, M., Young, V., Goldan, P. D., Kuster, W. C., Jobson, B. T., Fehsenfeld, F. C., Lonneman, W. A., Zika, R. D., Farmer, C. T., Riemer, D. D., and Rodgers, M. O.: Internal consistency tests for evaluation of measurements of anthropogenic hydrocarbons in the troposphere, J. Geophys. Res., 103, 22339–22359, 1998.

Paulson, S. E., Sen, A. D., Liu, P., Fenske, J. D., and Fox, M. J.: Evidence for formation of OH radicals from the reaction of O<sub>3</sub> with alkenes in the gas phase, Geophys. Res. Lett., 24, 3193–3196, 1997.

Platt, U., LeBras, G., Poulet, G., Burrows, J. P., and Moortgat, G.: Peroxy radicals from night-time reaction of NO<sub>3</sub> with organic compounds, Nature, 348, 147–149, 1990.

Platt, U., Alicke, B., Dubois, R., Geyer, A., Hofzumahaus, A., Holland F., Martinez, M., Mihelcic, D., Klöpfer, T., Lohrmann, B., Pätz, W., Perner, D., Rohrer, F., Schäfer, J., and Stutz, J.: Free radicals and fast photochemistry during BERLIOZ, J. Atmos. Chem., 42, 359–394, 2002.

Rivett, A. C., Martin, D., Nickless, G., Simmonds, P. G., O'Doherty, S. J., Gray, D. J., and Shallcross, D. E.: In-situ gas chromatographic measurements of halocarbons in an urban environment, Atmos. Environ., accepted, 2003.

Roberts, J. M., Fehsenfeld, F. C., Liu, S. C., Bollinger, M. J., Hahn, C., Albritton, D. L., and Sievers, R. E.: Measurements of aromatic hydrocarbon ratios and NO<sub>x</sub> concentrations in the rural troposphere: Observations of air mass photochemical ageing and NO<sub>x</sub> removal, Atmos. Environ., 18, 2421–2432, 1984.

Sheppard, V.: The near real time monitoring of NMHCs in tropospheric air. Thesis (Ph.D.), University of Bristol, 1999.

Sillman, S.: The relation between ozone, NO<sub>x</sub> and hydrocarbons in urban and polluted rural environments, Atmos. Environ., 33, 1821–1845, 1999.

Simmonds, P. G., O'Doherty, S. J., Nickless, G., Sturrock, G. A., Swaby, R., Knight, P., Ricketts, J., Woffendin, G., and Smith, R.: Automated gas-chromatograph mass-spectrometer for routine atmospheric measurements of the CFC replacement compounds, the hydrofluorocarbons and hydrochlorofluorocarbons, Anal. Chem., 67, 717–723, 1995.

Smith, N., Plane, J. M. C., Nien, C. F., and Solomon, P. A.: Nighttime radical chemistry in the San Joaquin valley, Atmos. Environ., 29, 2887–2897, 1995.

UK PORG: Ozone in the UK, UK Photochemical Oxidants Review Group Report, Department

ACPD

3, 769–796, 2003

## VOCs in the polluted urban environment of Bristol

A. C. Rivett et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003



of the Environment, London, 1997.

Wang, J. L., Chang, C. J., and Lin, Y. H.: Concentration distributions of anthropogenic halocarbons over a metropolitan area, Chemosphere, 36, 2391–2400, 1998.

Wayne, R. P.: Chemistry of Atmospheres (3rd Ed.), Oxford University Press, Oxford, 2000.

- 5 Yamamoto, N., Okayasu, H., Murayama, S., Mori, S., and Hunahashi, K.: Measurement of volatile organic compounds in the urban atmosphere of Yokohama, Japan, by an automated gas chromatographic system, Atmos. Environ., 34, 4441–4446, 2000.

**ACPD**

3, 769–796, 2003

---

**VOCs in the polluted  
urban environment of  
Bristol**

A. C. Rivett et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

**VOCs in the polluted  
urban environment of  
Bristol**

A. C. Rivett et al.


**Table 1.** Daytime hydroxyl radical concentrations during summer campaign (July to September), estimated from the mean ratios of four aromatic species to benzene

OH concentration (molecules cm <sup>-3</sup> )	
Average	$2.5 \times 10^6$
Maximum	$8.8 \times 10^6$
Minimum	$1.5 \times 10^5$

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

**VOCs in the polluted  
urban environment of  
Bristol**

A. C. Rivett et al.

**Fig. 1.** Time-series of halons and CFCs.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Print Version

Interactive Discussion

**VOCs in the polluted  
urban environment of  
Bristol**

A. C. Rivett et al.

**Fig. 2.** Time-series of 3 chlorinated compounds.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Print Version

Interactive Discussion

**VOCs in the polluted  
urban environment of  
Bristol**

A. C. Rivett et al.

**Fig. 3.** Time-series of TCE and PCE.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Print Version

Interactive Discussion

**VOCs in the polluted  
urban environment of  
Bristol**

A. C. Rivett et al.




**Fig. 4. (a)** Time-series of halons and CFCs overlaid with a representative hydrocarbon, toluene.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

**VOCs in the polluted  
urban environment of  
Bristol**

A. C. Rivett et al.




**Fig. 4. (b)** Time-series of the 3 chlorinated compounds overlaid with a representative hydrocarbon, toluene.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

**VOCs in the polluted  
urban environment of  
Bristol**

A. C. Rivett et al.




**Fig. 4. (c)** Time-series of TCE and PCE overlaid with a representative hydrocarbon, toluene.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)




**VOCs in the polluted  
urban environment of  
Bristol**

A. C. Rivett et al.

**Fig. 5.** Hydrocarbon time-series.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

**VOCs in the polluted  
urban environment of  
Bristol**

A. C. Rivett et al.




**Fig. 6.** Representative decay of hydrocarbons on the night of 30–31 August 2000.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

**VOCs in the polluted  
urban environment of  
Bristol**

A. C. Rivett et al.




**Fig. 7.** Steady-state night-time [OH] calculated for University (urban background) and NETCEN (urban centre) sites.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

**VOCs in the polluted  
urban environment of  
Bristol**

A. C. Rivett et al.




**Fig. 8. (a)** Percentage of total ozone formed, calculated by  $\varepsilon^{\text{POCP}}$  method.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

**VOCs in the polluted  
urban environment of  
Bristol**

A. C. Rivett et al.



**Fig. 8. (b)** Percentage of total ozone formed, calculated by  $\gamma_R$  method.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)